

Q1.

6 A compound, **A**, has the following composition by mass.

C, 66.7%; H, 11.1%; O, 22.2%.

It has an M_r of 72.

(a) Calculate the molecular formula of **A**.

Examiner's
Use

[2]

Q2.

2 Compounds of phosphorus have many uses in everyday life, e.g. fertilisers, matches and in water softeners.

(a) State the full electronic configuration of phosphorus.

.....[1]

(b) Phosphoric acid, H_3PO_4 , is used in the manufacture of phosphate fertilisers.

Deduce the oxidation number of phosphorus in H_3PO_4 .

.....[1]

(c) The salt sodium phosphate, Na_3PO_4 , is a water-softening agent.

(i) Write the equation for the complete neutralisation of phosphoric acid with aqueous sodium hydroxide.

.....

Sodium phosphate was prepared from 50.0 cm³ of 0.500 mol dm⁻³ H₃PO₄ and an excess of aqueous sodium hydroxide.

(ii) How many moles of H₃PO₄ were used?

(iii) Use your equation in (c)(i) to calculate how many moles of sodium hydroxide are required.

[3]

(d) Phosphorus sulphide, P₄S₃, is used in small amounts in the tip of a match. On striking a match, this compound burns.

(i) Construct an equation for this reaction.

.....

(ii) Both oxides formed in (i) dissolve in water to give acidic solutions. Construct an equation for the reaction of each oxide with water.

.....

.....[4]

[Total : 9]

Q3.

(c) 1.20 dm^3 of ammonia gas were dissolved in water to form 200 cm^3 of aqueous alkali at room temperature and pressure.

(i) Use the *Data Booklet* to calculate how many moles of $\text{NH}_3(\text{g})$ were dissolved.

(ii) Write the equation for the neutralisation of aqueous ammonia by dilute sulphuric acid.

.....

(iii) Calculate the volume of 0.50 mol dm^{-3} sulphuric acid that is required to neutralise the 200 cm^3 of aqueous ammonia.

[3]

Q4.

Hydrogen sulphide burns with a blue flame in an excess of oxygen to form sulphur dioxide and water.

Exa

(d) (i) Write a balanced equation for the complete combustion of H_2S .

.....

(ii) What is the change in the oxidation number of sulphur in this reaction?

from to

(iii) What volume of oxygen, measured at room temperature and pressure, is required for the complete combustion of 8.65 g of H_2S ? Give your answer to two decimal places.

[5]

Q5.

1 Mohr's salt is a pale green crystalline solid which is soluble in water. Mohr's salt is a 'double salt' which contains

Exa

two cations, one of which is Fe^{2+} ,

one anion which is SO_4^{2-} ,

and water of crystallisation.

(a) The identity of the second cation was determined by the following test. Solid Mohr's salt was heated with solid sodium hydroxide and a colourless gas was evolved. The gas readily dissolved in water giving an alkaline solution.

(i) What is the gas?

.....

(ii) What is the formula of the second cation identified by this test?

.....

(iii) In this test, a grey/green solid residue was also formed.

Suggest a name or formula for this solid.

.....

[3]

(b) The identity of the anion present in Mohr's salt was confirmed by adding dilute hydrochloric acid followed by aqueous barium chloride to an aqueous solution of Mohr's salt. A white precipitate was formed.

Suggest the identity of the white precipitate.

.....[1]

(c) When a double salt such as Mohr's salt is made, the two individual salts are mixed together in a 1:1 molar ratio, dissolved in water and the solution crystallised.

(i) Give the formula of **each** of the two salts that would be mixed to make the double salt, Mohr's salt.

salt 1

salt 2

(ii) Calculate the relative formula mass of **each** of the salts present in Mohr's salt.

salt 1

relative formula mass of salt 1

salt 2

relative formula mass of salt 2

(iii) The crystals of the double salt contain water of crystallisation.

The relative formula mass of Mohr's salt is 392. Use your answers to (ii) to calculate the number of moles of water of crystallisation present in one mole of Mohr's salt.

[6]

[Total: 10]

Q6.

- (c) In some countries, ethyne is manufactured from calcium carbide, CaC_2 , which is produced by heating quicklime and coke together at 2300 K.



When water is added to the CaC_2 , calcium hydroxide, Ca(OH)_2 , and ethyne, C_2H_2 , are produced.

- (i) Construct a balanced equation for the formation of ethyne from calcium carbide.

.....

- (ii) Use this equation and your answer to part (b) to calculate the mass of CaC_2 which will react with an excess of water to produce enough ethyne to fill 100 cylinders of the gas.

[3]

Q7.

- (b) When CH_2Cl_2 is heated under reflux with an excess of NaOH(aq) , a compound **W** is formed.

W has the following composition by mass: C, 40.0%; H, 6.7%; O, 53.3%.

Use this information and the *Data Booklet* to show that the empirical formula of **W** is CH_2O .

[2]

Q8.

Titanium also reacts with chlorine.

(d) When an excess of chlorine was reacted with 0.72 g of titanium, 2.85 g of a chloride **A** was formed.

(i) Calculate the amount, in moles, of titanium used.

(ii) Calculate the amount, in moles, of chlorine atoms that reacted.

(iii) Hence, determine the empirical formula of **A**.

(iv) Construct a balanced equation for the reaction between titanium and chlorine.

.....
[4]

Q9.

4 An organic compound, **E**, has the following composition by mass:
C, 48.7%; H, 8.1%; O, 43.2%.

(a) Calculate the empirical formula of **E**.

Ex

[2]

(b) When vaporised in a suitable apparatus, 0.130 g of **E** occupied a volume of 58.0 cm³ at 127 °C and $1.00 \times 10^5 \text{ Nm}^{-2}$.

(i) Use the expression $pV = \frac{mRT}{M_r}$ to calculate M_r of **E**,

where m is the mass of **E**.

(ii) Hence calculate the molecular formula of **E**.

[4]

Q10.

A third polycarboxylic acid present in unripe fruit is a colourless crystalline solid, **W**, which has the following composition by mass: C, 35.8%; H, 4.5%; O, 59.7%.


(d) (i) Show by calculation that the empirical formula of **W** is $\text{C}_4\text{H}_6\text{O}_5$.

(ii) The M_r of **W** is 134. Use this value to determine the molecular formula of **W**.

[3]

Q11.

- (c) In a redox reaction, 0.83 g of lithium reacted with water to form 0.50 dm³ of aqueous lithium hydroxide.



- (i) Calculate the amount, in moles, of lithium that reacted.

- (ii) Calculate the volume of hydrogen produced at room temperature and pressure.

- (iii) Calculate the concentration, in mol dm⁻³, of the LiOH(aq) formed.

[5]

Q12.

(c) In this section, give your answers to one decimal place.

The flight path from Beijing to Paris is approximately 8195 km.
A typical intercontinental jet airliner burns 10.8 kg of kerosene for each kilometre covered.

(i) Calculate the mass, in tonnes, of $C_{14}H_{30}$ burnt on a flight from Beijing to Paris.
[1 tonne = 1 000 kg]

(ii) Use your equation in **(b)** to calculate the mass, in tonnes, of CO_2 produced during this flight.

[4]

Q13.

In an experiment to determine K_c a student placed together in a conical flask 0.10 mol of ethanoic acid, 0.10 mol of an alcohol ROH, and 0.005 mol of hydrogen chloride catalyst. The flask was sealed and kept at 25 °C for seven days. After this time, the student titrated all of the contents of the flask with 2.00 mol dm⁻³ NaOH using phenolphthalein indicator. At the end-point, 22.5 cm³ of NaOH had been used.

- (b) (i) Calculate the amount, in moles, of NaOH used in the titration.
- (ii) What amount, in moles, of this NaOH reacted with the hydrogen chloride?
- (iii) Write a balanced equation for the reaction between ethanoic acid and NaOH.
- (iv) Hence calculate the amount, in moles, of NaOH that reacted with the ethanoic acid.

[4]

Q14.

- 1 Methanoic acid, HCO_2H , was formerly known as formic acid because it is present in the sting of ants and the Latin name for ant is *formica*. It was first isolated in 1671 by John Ray who collected a large number of dead ants and extracted the acid from them by distillation.

In this question, you should give all numerical answers to two significant figures.

At room temperature, pure methanoic acid is a liquid which is completely soluble in water.

When we are stung by a 'typical' ant a solution of methanoic acid, **A**, is injected into our skin.

Solution **A** contains 50% by volume of pure methanoic acid.

A 'typical' ant contains $7.5 \times 10^{-6} \text{ dm}^3$ of solution **A**.

(a) (i) Calculate the volume, in cm^3 , of solution **A** in one ant.

volume = cm^3

(ii) Use your answer to **(i)** to calculate the volume, in cm^3 , of pure methanoic acid in one ant.

volume = cm^3

(iii) Use your answer to **(ii)** to calculate how many ants would have to be distilled to produce 1 dm^3 of pure methanoic acid.

number =
[3]

When we are stung by an ant, the amount of solution **A** injected is 80% of the total amount of solution **A** present in one ant.

The density of pure methanoic acid is 1.2 g cm^{-3} .

(b) (i) Calculate the volume, in cm^3 , of **pure** methanoic acid injected in one ant sting.

volume = cm^3

(ii) Use your answer to **(i)** to calculate the mass of methanoic acid present in one ant sting.

mass = g
[3]

Bees also sting us by using methanoic acid. One simple treatment for ant or bee stings is to use sodium hydrogencarbonate, NaHCO_3 .

(c) (i) Construct a balanced equation for the reaction between methanoic acid and sodium hydrogencarbonate.

.....

(ii) In a typical bee sting, the mass of methanoic acid injected is $5.4 \times 10^{-3} \text{ g}$. Calculate the mass of NaHCO_3 needed to neutralise one bee sting.

mass = g
[3]

[Total: 9]

Q15.

2 Ammonium sulfate, $(\text{NH}_4)_2\text{SO}_4$, is widely used as a fertiliser.

E

In order to determine its percentage purity, a sample of ammonium sulfate fertiliser was analysed by reacting a known amount with an excess of $\text{NaOH}(\text{aq})$ and then titrating the unreacted NaOH with dilute HCl .

(a) Ammonium sulfate reacts with NaOH in a 1:2 ratio.
Complete and balance the equation for this reaction.



[2]

(b) A 5.00 g sample of a fertiliser containing $(\text{NH}_4)_2\text{SO}_4$ was warmed with 50.0 cm^3 (an excess) of $2.00 \text{ mol dm}^{-3} \text{ NaOH}$.

When all of the ammonia had been driven off, the solution was cooled.

The remaining NaOH was then titrated with $1.00 \text{ mol dm}^{-3} \text{ HCl}$ and 31.2 cm^3 were required for neutralisation.

(i) Write a balanced equation for the reaction between NaOH and HCl .

.....

(ii) Calculate the amount, in moles, of HCl in 31.2 cm^3 of $1.00 \text{ mol dm}^{-3} \text{ HCl}$.

(iv) Use your answers to (i), (ii) and (iii) to calculate the amount, in moles, of NaOH used up in the reaction with $(\text{NH}_4)_2\text{SO}_4$.

(v) Use your answer to (iv) and the equation in (a) to calculate the amount, in moles, of $(\text{NH}_4)_2\text{SO}_4$ that reacted with NaOH.

Exa

(vi) Use your answer to (v) to calculate the mass of $(\text{NH}_4)_2\text{SO}_4$ that reacted with NaOH.

(vii) Hence, calculate the percentage purity of the ammonium sulfate fertiliser.

[7]

[Total: 9]

Q16.

(e) The food additive E330 is another organic compound which occurs naturally in fruit. E330 has the following composition by mass: C, 37.5 %; H, 4.17 %; O, 58.3%. Calculate the empirical formula of E330.

[3]

Q17.

2 Washing soda is hydrated sodium carbonate, $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$.

E

A student wished to determine the value of x by carrying out a titration, with the following results.

5.13 g of washing soda crystals were dissolved in water and the solution was made up to 250 cm^3 in a standard volumetric flask.

25.0 cm^3 of this solution reacted exactly with 35.8 cm^3 of $0.100 \text{ mol dm}^{-3}$ hydrochloric acid and carbon dioxide was produced.

(a) (i) Write a balanced equation for the reaction between Na_2CO_3 and HCl .

.....

(ii) Calculate the amount, in moles, of HCl in the 35.8 cm^3 of solution used in the titration.

(iii) Use your answers to (i) and (ii) to calculate the amount, in moles, of Na_2CO_3 in the 25.0 cm^3 of solution used in the titration.

(iv) Use your answer to (iii) to calculate the amount, in moles, of Na_2CO_3 in the 250 cm^3 of solution in the standard volumetric flask.

(v) Hence calculate the mass of Na_2CO_3 present in 5.13 g of washing soda crystals.

Ex

[6]

(b) Use your calculations in (a) to determine the value of x in $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$.

[2]

[Total: 8]

Q18.

1 A sample of a fertiliser was known to contain ammonium sulfate, $(\text{NH}_4)_2\text{SO}_4$, and sand only.

A 2.96 g sample of the solid fertiliser was heated with 40.0 cm^3 of $\text{NaOH}(\text{aq})$, an excess, and all of the ammonia produced was boiled away.

After cooling, the remaining $\text{NaOH}(\text{aq})$ was exactly neutralised by 29.5 cm^3 of 2.00 mol dm^{-3} HCl .

In a separate experiment, 40.0 cm^3 of the original $\text{NaOH}(\text{aq})$ was exactly neutralised by 39.2 cm^3 of the 2.00 mol dm^{-3} HCl .

In a separate experiment, 40.0 cm^3 of the original NaOH(aq) was exactly neutralised by 39.2 cm^3 of the $2.00 \text{ mol dm}^{-3} \text{ HCl}$.

(a) (i) Write balanced equations for the following reactions.

NaOH with HCl

.....

$(\text{NH}_4)_2\text{SO}_4$ with NaOH

.....

(ii) Calculate the amount, in moles, of NaOH present in the 40.0 cm^3 of the original NaOH(aq) that was neutralised by 39.2 cm^3 of $2.00 \text{ mol dm}^{-3} \text{ HCl}$.

(iii) Calculate the amount, in moles, of NaOH present in the 40.0 cm^3 of NaOH(aq) that remained after boiling the $(\text{NH}_4)_2\text{SO}_4$.

(iv) Use your answers to **(ii)** and **(iii)** to calculate the amount, in moles, of NaOH that reacted with the $(\text{NH}_4)_2\text{SO}_4$.

(v) Use your answers to (i) and (iv) to calculate the amount, in moles, of $(\text{NH}_4)_2\text{SO}_4$ that reacted with the NaOH.

(vi) Hence calculate the mass of $(\text{NH}_4)_2\text{SO}_4$ that reacted.

(vii) Use your answer to (vi) to calculate the percentage, by mass, of $(\text{NH}_4)_2\text{SO}_4$ present in the fertiliser.

Write your answer to a suitable number of significant figures.

[9]

(b) The uncontrolled use of nitrogenous fertilisers can cause environmental damage to lakes and streams. This is known as *eutrophication*.

What are the processes that occur when excessive amounts of nitrogenous fertilisers get into lakes and streams?

.....
.....
..... [2]

(c) Large quantities of ammonia are manufactured by the Haber process.

Not all of this ammonia is used to make fertilisers.

State **one** large-scale use for ammonia, **other than** in the production of nitrogenous fertilisers.

..... [1]

[Total: 12]


Q19.

- 2 Chile saltpetre is a mineral found in Chile and Peru, and which mainly consists of sodium nitrate, NaNO_3 . The mineral is purified to concentrate the NaNO_3 which is used as a fertiliser and in some fireworks.


In order to find the purity of a sample of sodium nitrate, the compound is heated in $\text{NaOH}(\text{aq})$ with Devarda's alloy which contains aluminium. This reduces the sodium nitrate to ammonia which is boiled off and then dissolved in acid.



The ammonia gas produced is dissolved in an excess of H_2SO_4 of known concentration.



The amount of unreacted H_2SO_4 is then determined by back-titration with NaOH of known concentration.



- (a) A 1.64 g sample of impure NaNO_3 was reacted with an excess of Devarda's alloy. The NH_3 produced was dissolved in 25.0 cm^3 of $1.00 \text{ mol dm}^{-3} \text{ H}_2\text{SO}_4$. When all of the NH_3 had dissolved, the resulting solution was titrated with $\text{NaOH}(\text{aq})$. For neutralisation, 16.2 cm^3 of $2.00 \text{ mol dm}^{-3} \text{ NaOH}$ were required.
- (i) Calculate the amount, in moles, of H_2SO_4 present in the 25.0 cm^3 of $1.00 \text{ mol dm}^{-3} \text{ H}_2\text{SO}_4$.
- (ii) Calculate the amount, in moles, of NaOH present in 16.2 cm^3 of $2.00 \text{ mol dm}^{-3} \text{ NaOH}$.
- (iii) Use your answer to (ii) to calculate the amount, in moles, of H_2SO_4 that reacted with 16.2 cm^3 of $2.00 \text{ mol dm}^{-3} \text{ NaOH}$.

(iv) Use your answers to (i) and (iii) to calculate the amount, in moles, of H_2SO_4 that reacted with the NH_3 .

(v) Use your answer to (iv) to calculate the amount, in moles, of NH_3 that reacted with the H_2SO_4 .

(vi) Use your answer to (v) to calculate the amount, in moles, of NaNO_3 that reacted with the Devarda's alloy.

(vii) Hence calculate the mass of NaNO_3 that reacted.

(viii) Use your answer to (vii) to calculate the percentage by mass of NaNO_3 present in the impure sample.
Write your answer to a suitable number of significant figures.

[9]

(b) The above reaction is an example of a redox reaction.
What are the oxidation numbers of nitrogen in NaNO_3 and in NH_3 ?

NaNO_3

NH_3

[1]

[Total: 10]

Q20.

- 1 Most submarines travel under water using electrical power from batteries. The German engineer Helmut Walter designed a diesel engine that could be used to propel a submarine beneath the surface of the sea. Instead of taking air from above the surface of the sea, Walter's engine used hydrogen peroxide, H_2O_2 , to provide oxygen for a conventional diesel engine.

Hydrogen peroxide may be catalytically decomposed to give water and oxygen.


- (a) (i) What is meant by the term *catalyst*?

.....
.....

- (ii) Construct a balanced equation for the decomposition of H_2O_2 .

..... [3]

Diesel fuel may be considered to consist of the hydrocarbon $\text{C}_{15}\text{H}_{32}$ which reacts completely with oxygen according to the following equation.



- (b) (i) To which homologous series does $\text{C}_{15}\text{H}_{32}$ belong?

.....

- (ii) Use the equation above and your answer to (a)(ii) to calculate the amount, in moles, of H_2O_2 , that will provide sufficient oxygen for the complete oxidation of one mole of $\text{C}_{15}\text{H}_{32}$.

amount of H_2O_2 = mol

[3]

A submarine equipped with a Walter engine used 212 tonnes of diesel fuel during an underwater voyage. The submarine also carried concentrated aqueous H_2O_2 .
[1 tonne = 10^6 g]

For
Examiner's
Use

(c) (i) Calculate the amount, in moles, of diesel fuel used during the underwater voyage.

amount of diesel fuel = mol

(ii) Use your answers to (b)(ii) and (c)(i) to calculate the mass, in tonnes, of hydrogen peroxide used during the underwater voyage.

mass of H_2O_2 = tonnes
[4]

(d) The exhaust products of the Walter engine were passed into the sea.

What would happen to them?

..... [1]

[Total: 11]

Q21.


1 In 1814, Sir Humphrey Davy and Michael Faraday collected samples of a flammable gas, **A**, from the ground near Florence in Italy. They analysed **A** which they found to be a hydrocarbon. Further experiments were then carried out to determine the molecular formula of **A**.

(a) What is meant by the term *molecular formula*?

.....
.....
..... [2]

Davy and Faraday deduced the formula of **A** by exploding it with an excess of oxygen and analysing the products of combustion.

- (b) Complete and balance the following equation for the complete combustion of a hydrocarbon with the formula C_xH_y .



- (c) When 10 cm^3 of **A** was mixed at room temperature with 50 cm^3 of oxygen (an excess) and exploded, 40 cm^3 of gas remained after cooling the apparatus to room temperature and pressure.
When this 40 cm^3 of gas was shaken with an excess of aqueous potassium hydroxide, KOH, 30 cm^3 of gas still remained.

- (i) What is the identity of the 30 cm^3 of gas that remained at the end of the experiment?

.....

- (ii) The combustion of **A** produced a gas that reacted with the KOH(aq).

What is the identity of this gas?

.....

- (iii) What volume of the gas you have identified in (ii) was produced by the combustion of **A**?

..... cm^3

- (iv) What volume of oxygen was used up in the combustion of **A**?

..... cm^3

[4]

- (d) Use your equation in (b) and your results from (c)(iii) and (c)(iv) to calculate the molecular formula of **A**.
Show all of your working.

[3]

[Total: 11]

Q22.

- 1 Compound **A** is an organic compound which contains carbon, hydrogen and oxygen.

When 0.240 g of the vapour of **A** is slowly passed over a large quantity of heated copper(II) oxide, CuO, the organic compound **A** is completely oxidised to carbon dioxide and water. Copper is the only other product of the reaction.

The products are collected and it is found that 0.352 g of CO₂ and 0.144 g of H₂O are formed.

- (a) In this section, give your answers to three decimal places.

- (i) Calculate the mass of carbon present in 0.352 g of CO₂.

Use this value to calculate the amount, in moles, of carbon atoms present in 0.240 g of **A**.

(ii) Calculate the mass of hydrogen present in 0.144 g of H₂O.

Use this value to calculate the amount, in moles, of hydrogen atoms present in 0.240 g of **A**.

(iii) Use your answers to calculate the mass of oxygen present in 0.240 g of **A**.

Use this value to calculate the amount, in moles, of oxygen atoms present in 0.240 g of **A**.

[6]

(b) Use your answers to (a) to calculate the empirical formula of **A**.

[1]

(c) When a 0.148 g sample of **A** was vapourised at 60°C, the vapour occupied a volume of 67.7 cm³ at a pressure of 101 kPa.

(i) Use the general gas equation $pV = nRT$ to calculate M_r of **A**.

$M_r = \dots\dots\dots$

(ii) Hence calculate the molecular formula of **A**.

[3]

Q23.

2 When 0.42 g of a gaseous hydrocarbon **A** is slowly passed over a large quantity of heated copper(II) oxide, CuO, **A** is completely oxidised.

The products are collected and it is found that 1.32 g of CO₂ and 0.54 g of H₂O are formed. Copper is the only other product of the reaction.

(a) (i) Calculate the mass of carbon present in 1.32 g of CO₂.

Use this value to calculate the amount, in moles, of carbon atoms present in 0.42 g of **A**.

(ii) Calculate the mass of hydrogen present in 0.54 g of H₂O.

Use this value to calculate the amount, in moles, of hydrogen atoms present in 0.42 g of **A**.

(iii) It is thought that **A** is an alkene rather than an alkane.

Use your answers to (i) and (ii) to deduce whether this is correct.

Explain your answer.

.....

..... [5]

(b) Analysis of another organic compound, **B**, gave the following composition by mass: C, 64.86%; H, 13.50%, O, 21.64%.

E

(i) Use these values to calculate the empirical formula of **B**.

Q24.

1 Zinc is an essential trace element which is necessary for the healthy growth of animals and plants. Zinc deficiency in humans can be easily treated by using zinc salts as dietary supplements.

(a) One salt which is used as a dietary supplement is a hydrated zinc sulfate, $\text{ZnSO}_4 \cdot x\text{H}_2\text{O}$, which is a colourless crystalline solid.

Crystals of zinc sulfate may be prepared in a school or college laboratory by reacting dilute sulfuric acid with a suitable compound of zinc.

Give the formulae of **two** simple compounds of zinc that could **each** react with dilute sulfuric acid to produce zinc sulfate.

..... and [2]

(b) A simple experiment to determine the value of x in the formula $\text{ZnSO}_4 \cdot x\text{H}_2\text{O}$ is to heat it carefully to drive off the water.



A student placed a sample of the hydrated zinc sulfate in a weighed boiling tube and reweighed it. He then heated the tube for a short time, cooled it and reweighed it when cool. This process was repeated four times. The final results are shown below.

mass of empty tube /g	mass of tube + hydrated salt /g	mass of tube + salt after fourth heating /g
74.25	77.97	76.34

(i) Why was the boiling tube heated, cooled and reweighed four times?

.....
.....

(ii) Calculate the amount, **in moles**, of the anhydrous salt produced.

(iii) Calculate the amount, **in moles**, of water driven off by heating.

(iv) Use your results to (ii) and (iii) to calculate the value of x in $\text{ZnSO}_4 \cdot x\text{H}_2\text{O}$.

[7]

(c) For many people, an intake of approximately 15 mg per day of zinc will be sufficient to prevent deficiencies.

Zinc ethanoate crystals, $(\text{CH}_3\text{CO}_2)_2\text{Zn} \cdot 2\text{H}_2\text{O}$, may be used in this way.

(i) What mass of pure crystalline zinc ethanoate ($M_r = 219.4$) will need to be taken to obtain a dose of 15 mg of zinc?

(ii) If this dose is taken in solution as 5 cm^3 of aqueous zinc ethanoate, what would be the concentration of the solution used?
Give your answer in mol dm^{-3} .

[4]

[Total: 13]

Q25.

- 1 Carbon dioxide, CO_2 , makes up about 0.040% of the Earth's atmosphere. It is produced by animal respiration and by the combustion of fossil fuels.

In animal respiration, oxygen reacts with a carbohydrate such as glucose to give water, carbon dioxide and energy.

The typical daily food requirement of a human can be considered to be the equivalent of 1.20 kg of glucose, $\text{C}_6\text{H}_{12}\text{O}_6$.

You should express all of your numerical answers in this question to three significant figures.

- (a) (i) Construct a balanced equation for the complete oxidation of glucose.

.....

- (ii) Use your equation to calculate the amount, in moles, of CO_2 produced by one person in one day from 1.20 kg of glucose.

- (iii) On the day on which this question was written, the World population was estimated to be 6.82×10^9 .

Calculate the total mass of CO_2 produced by this number of people in one day. Give your answer in tonnes. [1 tonne = 1.00×10^6 g]

[5]

- (b)** When fossil fuels are burned in order to give energy, carbon dioxide and water are also produced.

Exa
t

The hydrocarbon octane, C_8H_{18} , can be used to represent the fuel burned in motor cars. A typical fuel-efficient motor car uses about 4.00 dm^3 of fuel to travel 100 km.

- (i)** Construct a balanced equation for the complete combustion of octane.

.....

- (ii)** The density of octane is 0.700 g cm^{-3} .

Calculate the amount, in moles, of octane present in 4.00 dm^3 of octane.

- (iii)** Calculate the mass of CO_2 produced when the fuel-efficient car is driven for a distance of 100 km.

[5]

- (c)** Calculate how many kilometres the same fuel-efficient car would have to travel in order to produce as much CO_2 as is produced by the respiration of 6.82×10^9 people during one day. Use your answer to **(a)(iii)**.

[2]

- (d) Carbon dioxide is one of a number of gases that are responsible for global warming. When fossil fuels such as octane are burned in a car engine, other atmospheric pollutants are also produced. Give the formula of **one** atmospheric pollutant that may be produced in a car engine, other than CO_2 , and state how this pollutant damages the environment.

pollutant

damage caused

[2]

[Total: 14]

Q26.

4 Compound **R** is a weak diprotic (dibasic) acid which is very soluble in water.

*For
Examiner's
Use*

- (a) A solution of **R** was prepared which contained 1.25 g of **R** in 250 cm^3 of solution. When 25.0 cm^3 of this solution was titrated with $0.100 \text{ mol dm}^{-3}$ NaOH, 21.6 cm^3 of the alkali were needed for complete reaction.

- (i) Using the formula H_2X to represent **R**, construct a balanced equation for the reaction between H_2X and NaOH.

.....

- (ii) Use the data above to calculate the amount, in moles, of OH^- ions used in the titration.

- (iii) Use your answers to (i) and (ii) to calculate the amount, in moles, of **R** present in 25.0 cm^3 of solution.

- (iv) Calculate the amount, in moles, of **R** present in 250 cm^3 of solution.

.....

(v) Calculate M_r of **R**.

[5]

(b) Three possible structures for **R** are shown below.

S	T	U
$\text{HO}_2\text{CCH}=\text{CHCO}_2\text{H}$	$\text{HO}_2\text{CCH}(\text{OH})\text{CH}_2\text{CO}_2\text{H}$	$\text{HO}_2\text{CCH}(\text{OH})\text{CH}(\text{OH})\text{CO}_2\text{H}$

(i) Calculate the M_r of each of these acids.

M_r of **S** = M_r of **T** = M_r of **U** =

(ii) Deduce which of the structures, **S**, **T** or **U**, correctly represents the structure of the acid, **R**.

R is represented by

[2]

